



GB 04/442

INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

# **PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 17 NOV 2004


WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




Signed

Dated

*R. Mahoney*  
9 November 2004

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977  
(Rule 16)21 OCT 03 EB46175-1 0324523.0  
P01/7700 0.00-0324523.0

The Patent Office

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form.)

Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your reference

ARDW / P28582GB

2. Patent application number

(The Patent Office will fill in this part)

0324523.0

21 OCT 2003

3. Full name, address and postcode of the or of each applicant (underline all surnames)Medical Research Council  
20 Park Crescent  
London  
W1B 1AL  
United Kingdom

0324523.0

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

59607001

4. Title of the invention

COMPOSITIONS AND METHODS OF TREATMENT

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

ERIC POTTER CLARKSON  
PARK VIEW HOUSE  
58 THE ROPEWALK  
NOTTINGHAM  
NG1 5DD

Patents ADP number (if you know it)

1305010

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

YES

- a) any applicant named in part 3 is not an inventor; or
- b) there is an inventor who is not named as an applicant; or
- c) any named applicant is a corporate body.  
See note (d))

Patents Form 1/77

## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 56

Claims(s) 11

Abstract 1

Drawing(s) 16 only

10. If you are also filing in any of the following, state how many against each item.

Priority Documents 0

Translations of priority documents 0

Statement of inventorship and right to grant of a patent (Patents Form 7/77) NO

Request for preliminary examination and search (Patents Form 9/77) YES

Request for substantive examination (Patents Form 10/77) NO

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

ERIC POTTER CLARKSON

Date

21 October 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

0115 9552211

### Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 01645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

DUPLICATE

1

**COMPOSITIONS AND METHODS OF TREATMENT**

The present invention relates to therapeutic compositions, methods and uses; in particular it relates to methods for treating degenerative diseases in a patient.

Many serious medical conditions, such as Type I diabetes, osteoarthritis, rheumatoid arthritis, multiple sclerosis, heart failure, stroke, burns, osteoporosis, bone fractures, Parkinson's disease and spinal chord injury, are due to the failure of or damage to tissue or a cell type within a patient due to disease or trauma. These can be considered degenerative diseases, some of which are associated with aging and where cells are unable to repair themselves or be replaced. Current treatments are limited to being palliative, delaying progression, and tissue function is typically not restored. Recent breakthroughs in the isolation, expansion and controlled differentiation of human adult and embryonic stem cells and the restoration of normal tissue function in animal models of degenerative disease following experimental transplantation have opened up the possibility of a new major field of regenerative medicine. New procedures are being developed to correct the failure of or damage to the tissue or cell type concerned by introducing into the patient cells which are able to take the place and function of the failed or damaged tissue or cells. In some cases, these may be exactly the same type of cells that are failed or damaged. In other cases, the introduced cells will be precursor cells of the tissue or cell type to be replaced, which are able to differentiate into the desired tissue or cell type at the site of disease or injury. In some cases, cells will be introduced at the precise site of disease or trauma; in others, cells will be introduced into portal veins, ventricles or elsewhere in the vasculature, circulatory or lymphatic systems to facilitate migration to the site of disease or trauma.

30



An example of this approach is in relation to Parkinson's disease, which is a very common neurodegenerative disorder that affects more than 2% of the population over 65 years of age. Parkinson's disease is caused by a progressive degeneration and loss of dopamine-producing neurons, which leads to tremor, rigidity and hypokinesia (abnormally decreased mobility). A recent study has shown that mouse embryonic stem cells can differentiate into dopamine-producing neurons by introducing the Nurr1 gene. When transplanted into the brains of a rat model of Parkinson's disease, these stem cell-derived dopamine-producing neurons reinnervated the brains of the rat Parkinson models, released dopamine and improved motor function.

~~A further example of the approach is the use of cardiomyocytes or bone marrow stem cells to repair damage to heart muscle tissue for example in chronic heart disease or after an infarction. A still further example is the use of oligodendrocytes for repairing damage to the spinal chord. A yet still further example is the use of derivatives of human embryonic stem cells which are able to differentiate into insulin-producing cells that can be used in transplantation therapy to treat Type I diabetes.~~

Useful information on stem cells and their use in regenerative medicine may be found on the National Institutes of Health web site, for example at <http://stemcells.nih.gov>. In addition, the potential of stem cells is reviewed by Pfendler & Kawase (2003) *Obstetrical & Gynecological Survey* 58, 197-208, incorporated herein by reference.

One of the most important applications of human stem cells, therefore, is the generation of cells and tissues that can be used for cell-based therapies. Today, donated organs and tissues are often used to replace failing or destroyed tissue, but the need for transplantable tissues and organs far

outweighs the available supply, hence the great interest in cell-based, particularly stem cell-based therapy.

To realise the potential of cell-based therapies for such pervasive and debilitating diseases as those discussed above, it is necessary for the cells to survive in the patient after transplantation. Unless the cell used in the therapy, such as the stem cell, is derived from the patient, it is highly likely that the patient will raise an immune response to it, thereby reducing the chances of it being rejected by the patient's immune system, and increasing the likelihood that immunosuppressive drugs will have to be used. This is because the introduced cell is considered to be "foreign" by the patient's immune system because of the presence of "foreign" antigens on the cell. Indeed, this is the conclusion reached in relation to human embryonic stem cells, where Drukker *et al* (2002) *Proc. Natl. Acad. Sci. USA* 99, 9864-9869 notes that these cells can express high levels of MHC-I proteins and thus may be rejected on transplantation. However, in Bell (2002) *Nature Reviews Immunology* 2, 75 there is a suggestion that embryonic stem cells may survive in allogeneic hosts in the absence of host conditioning.

Ways to reduce the possibility of undesirable immune responses and rejection of cells used in therapy have been suggested. For example, a "master" embryonic stem cell line may be produced in which the major histocompatibility complex (MHC) genes have been genetically modifying or knocked out. However, this may be technically difficult to achieve and, if accomplished, could expose the recipient of the MHC null transplant to new risks of infectious disease and/or cancer. An alternative strategy that has been suggested is to introduce the recipient's MHC genes into the embryonic stem cell through targeted gene transfer, but because of the differences among MHC proteins among individuals, the donor stem cells may be recognised as non-self by the patient's immune system and trigger graft versus host disease

(ie destruction by cytotoxic T cells) and ultimately rejection. Furthermore, this approach is also technically demanding and complex.

An organism's immunity to an antigen arises as a consequence of a first encounter with the antigen and the subsequent production of immunoglobulin molecules, for example, antibodies, capable of selectively binding that antigen. In addition, the immune response is controlled by T cells which may be antigen specific. A large proportion of the memory T-cell population (8-10%) will recognise MHC antigens. Immunity allows the rapid recruitment, usually by stimulating an inflammatory response, of cells which can dispose of the foreign antigen. Under certain circumstances, the immune system does not produce an immune response against antigens due to a mechanism called "tolerance". For example, an immune system can normally discriminate against foreign antigens and constituents of the organism itself, due to a mechanism whereby all T and B lymphocytes which could potentially produce antibodies to constituents of the organism itself ("self antigens") are destroyed during development, thereby removing the organism's capacity to produce antibodies directed to a self antigen.

One way that has been suggested of tolerising a patient who is undergoing a cell transplant is to have pre-tolerised the patient to the MHC antigens of the "master" embryonic stem cell line from which the cell or tissue for transplantation will be derived. This requires a procedure somewhat akin to a bone marrow transplant, and so certainly is invasive and requires some degree of immunosuppression.

The inventors now describe a much simpler method for inducing tolerance in (or "pre-tolerising") a patient to a cell or tissue which "regenerates" failed or damaged cells or tissues in the patient by producing a tolerant environment in the patient into which a cell is introduced which is a precursor of the cell or

tissue to be generated. In particular, the tolerant environment into which the precursor cell is introduced is created using an agent which is able to raise the effective cAMP concentration in a monocyte cell, such as a prostaglandin, preferably in combination with granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof. Typically, the prostaglandin may also be used in combination with a phosphodiesterase inhibitor.

It has been found that there is a synergistic effect between prostaglandin and a phosphodiesterase (PDE) inhibitor on the release of interlenkin-10 (IL-10) from cells of the immune system. Furthermore, it has been found that there is a marked stimulation of IL-10 and inhibition of interleukin-12 (IL-12) in cells of the immune system when a prostaglandin and a PDE inhibitor are used in combination. In the presence of a PDE inhibitor, the stimulation of IL-10 by both PGE and 19-hydroxy PGE was increased strikingly, resulting in a tolerising environment.

PDE inhibitors such as Rolipram are known to raise cAMP and IL-10 levels in monocyte/macrophages stimulated with the bacterial coat product lipopolysaccharide (LPS) (Strassman *et al* (1994) *J. Exp. Med.* 180: 2365-70; Kraan *et al* (1995) *J. Exp. Med.* 181: 775-9; Kambayashi *et al* (1995) *J. Immunol.* 155: 4909-16).

It has also been shown that an increase in PDE activity follows both PGE and 19-hydroxy PGE application. This is a direct negative feedback to reduce the effect of the stimulus. Use of a PGE and a PDE inhibitor increases PDE message even further, but then the synthesised phosphodiesterase is nullified by the presence of the inhibitor.

The principal receptors for prostaglandin E2 (PGE2) are the EP2 and EP4 subtypes; however, other receptor sub-types exist (namely EP1 and EP3). EP2

and EP4 receptors couple with adenylylase and use elevated cAMP as the messenger system. The levels of cAMP in tissue are governed both by synthesis and by catabolism by PDE. PDE can be blocked by specific inhibitors. The inventors believe, but without being bound by any theory, that the administration of a PDE inhibitor will enhance the effect of a prostaglandin or agonist thereof in inducing tolerance to a precursor cell (or an antigen found thereon or a derivative thereof) that is administered to a patient. Thus, the inventors believe, but without being bound by any theory, that the effect of a prostaglandin or agonist thereof (such as PGE) acting on its EP2 and EP4 receptors is to stimulate cAMP and the addition of the PDE inhibitor provides a synergistic action on monocytes and macrophages resulting in a reduction in the immune and/or inflammatory response which is greater than the effect of the sum of the same amount of either prostaglandin or agonist thereof or PDE inhibitor administered alone.

15

It has also been found that there is a marked stimulation of IL-10 in cells of the immune system when an agent which raises the effective cAMP concentration in monocyte cells, such as a prostaglandin, and granulocyte-macrophage colony stimulating factor (GMCSF) are used in combination. It has been found that there is a synergistic effect between a prostaglandin and GMCSF on the release of IL-10 from cells of the immune system; in the presence of GMCSF the stimulation of IL-10 by both prostaglandin E (PGE) and 19-hydroxy PGE was increased strikingly, resulting in a tolerising environment. In other words, it is believed that GMCSF and an agent that raises the effective cAMP concentration in a monocyte cell, such as a prostaglandin, polarises monocytes into a phenotype characterised by increased IL-10 release. Similarly, in the presence of GMCSF the stimulation of IL-10 expression by forskolin is increased strikingly, and in a synergistic way compared to forskolin or GMCSF alone. Not only is the cell directed to a pro-tolerance phenotype but this is also accompanied by enhanced production

30

of granulysin, an anti-microbial agent. In addition, the effects of PGE and GMCSF are prolonged and continue after the removal of these agents thus the cell is selectively differentiated.

- 5 GMCSF has an important role in granulocyte and macrophage lineage maturation. GMCSF has been proposed as both a treatment agent and a target for treatment. Recombinant human GMCSF has been used to treat some cancers and to promote haematopoietic reconstitution following bone marrow transplantation (Leukine<sup>®</sup> Package Insert Approved Text, February 1998, and
- 10 Buchsel *et al* (2002) *Clin. J. Oncol. Nurs.* 6(4): 198-205). By contrast, other recent reports describe GMCSF as being a potential target for treatment of inflammatory and immune diseases (Hamilton (2002) *Trends Immunol* 23(8): 403-8) and asthma (Ritz *et al* (2002) *Trends Immunol* 23(8): 396-402).
- 15 In diseases resulting from an aberrant or undesired immune response there is often a deficiency in IL-10. This deficiency in IL-10 may be detrimental to the development of useful T helper cells, particularly type-2 T helper cells; a preponderance of type 1 T helper cells over type 2 T helper cells is thought to be characteristic of autoimmune disease. Thus, stimulation of IL-10
- 20 production is believed to induce a tolerising environment for T cell priming. In addition, a high IL-10 environment will act on an antigen presenting cell (typically a dendritic cell) to ensure regulatory T cell formation, creating a regulatory T cell that is specific for the antigen presented.
- 25 Without being bound by theory, the inventor believes that a combination of GMCSF and an agent which raises the effective cAMP concentration in a monocyte cell, such as a prostaglandin or forskolin, will also decrease IL-12 levels, which would be expected to enhance the effects of the invention. It has been shown that the combination of a prostaglandin and GMCSF increases the
- 30 expression of both IL-10 and COX-2, and that the combination of a forskolin

and GMCSF synergistically increases the level of IL-10 in a monocyte cell. The decrease in IL-12 levels may therefore arise through the direct inhibition of IL-12 by IL-10 (Harizi *et al* (2002) *J. Immunol.* **168**, 2255-2263) or through an IL-10 independent pathway that depends on COX-2 induction (Schwacha *et al* (2002) *Am. J. Physiol. Cell Physiol.* **282**, C263-270).

It has also been shown that PGE and GMCSF reduce levels of participants in antigen presentation such as class II transactivator (CIITA) and MHC class II (as shown in Example 1). This change in phenotype is accompanied by enhanced expression of granulysin which has antimicrobial, including antiviral, properties (Krensky (2000) *Biochem. Pharmacol.* **59**, 317-320) and is normally thought of as a product of activated T-cells that mediates antiviral activity that lyses infected cells (Hata *et al* (2001) *Viral Immunol.* **14**, 125-133; Ochoa *et al* (2001) *Nature Medicine* **7**, 174-179; Smyth *et al* (2001) *J. Leukoc. Biol.* **70**, 18-29). The increased expression of granulysin is believed to be an important consequence of the present invention, as the increase in innate defence molecules may compensate for the compromise of the adaptive immune system that accompanies tolerance induction.

In addition, it has been shown that a combination of PGE and GMCSF increases the expression of COX-2, CD86, CD14. COX-2 is believed to be involved in maintaining the tolerant phenotype after removal of the prostaglandin and GMCSF (as is shown in Examples 2 and 3), and both CD14 and CD86 are differentiation markers and are evidence of a more differentiated state.

The inventors now propose inducing tolerance to a cell in a patient by the use of an agent which raises the effective cAMP concentration in a monocyte cell in order to induce a tolerising environment in the patient, and by administering the cell or a precursor thereof or an antigen found thereon or a derivative of

said antigen to the patient, such that tolerance to the cell is induced in the patient. By this process the patient is also made tolerant to a therapeutic cell which has the same antigenic characteristics as the cell used for tolerisation.

- 5 As far as the inventors are aware, no-one has suggested the use of this system of generating tolerance in connection with cell based therapies or its use in cellular transplants for treating degenerative disease.

- 10 The listing or discussion of a prior-published document in this specification should not necessarily be taken as an acknowledgement that the document is part of the state of the art or is common general knowledge.

- 15 A first aspect of the invention provides a method of inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof, the method comprising administering to the patient (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.

- 20 By inducing tolerance, we include the meaning that when the patient is subsequently administered the therapeutic cell or a precursor thereof, a greatly reduced or non-damaging immune response with respect to the therapeutic cell is experienced by the patient compared to a patient who has not been pretolerised. The well known mixed lymphocyte test may be used to  
25 determine whether a patient has been pretolerised. Alternatively, loss of the 6C10 marker (as described in Maier *et al* (1998) *Proc. Natl. Acad. Sci. USA* 95, 4499) may be used.



It will be appreciated that inducing tolerance in, or pretolerising, the patient is beneficial in those patients who will subsequently be administered a therapeutic amount of the therapeutic cells when undergoing transplantation for the purpose of repairing or regenerating damaged cells or tissue. A  
5 therapeutic amount of the cells is the amount which is needed to be administered to the patient in order to achieve a beneficial effect in terms of satisfying the need of the patient, for example in combating a degenerative disease or disorder. The therapeutic cells typically are used to repair or regenerate failed or damaged cells or tissues.

10

It will be appreciated that a benefit of inducing tolerance (or pretolerising) is that the chances of an adverse reaction on subsequent transplantation of the therapeutic cells is reduced.

15 By "sharing the same antigenic characteristics" we include the meaning that the tolerising cell has sufficient cell surface antigens, typically MHC antigens, in common with the therapeutic cell that administration of the tolerising cell to the patient in a tolerising environment leads to tolerance to the subsequent administration of the therapeutic cell so that the likelihood of rejection of the  
20 therapeutic cell is reduced compared to when no tolerisation is used.

Typically, the tolerising cell and therapeutic cell are syngeneic and express substantially the same antigens.

25 A second aspect of the invention provides a method of reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration, the method comprising administering to the patient prior to the transplant (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell which therapeutic cell is, or is able to  
30 differentiate into, the cell or tissue to be regenerated, or an antigen found

thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.

5 A third aspect of the invention provides a method of treating a patient in need of cell or tissue regeneration the method comprising administering to the patient (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the  
10 effective cAMP concentration in a monocyte cell in an amount to induce tolerance to the said therapeutic cell, and subsequently administering to the patient (c) a therapeutic amount of the said therapeutic cell.

15 Although the invention may be used in connection with any mammal, including domestic and farm mammals such as cat, dog, horse, sheep, cow and the like, typically the patient is a human.

The diseases where the methods of the invention may be used include degenerative diseases or disorders by which we include diabetes, where  
20 insulin-producing cells fail; stroke, Parkinson's disease, ALS (Lou Gehrig's disease) and spinal cord injury, where nerve cells fail; heart attack, cardiac ischaemia and congestive heart failure, where heart muscle cells fail; cirrhosis and hepatitis, where liver cells fail; certain cancers and immunodeficiency, where blood, bone marrow or haematopoietic cells fail; osteoporosis, where  
25 bone cells fail; osteoarthritis, where cartilage cells fail; burns and wounds, where skin cells fail; muscular dystrophy, where skeletal muscle cells fail; age-related macular degeneration where retinal cells fail; and multiple sclerosis, where myelin is destroyed (Schwann cells fail).

It will be appreciated that in preferred embodiments of the practice of the methods of the invention there are two distinct points at which the patient is administered a cell. The first is in the context of tolerising the patient, and the second is in the context of administering a cell in therapeutic amounts once the  
5 patient has been tolerised to the cell.

The first cell ("tolerising cell") may be any suitable cell which shares the same antigenic characteristics as the second cell ("therapeutic cell"). Thus, typically and preferably, the tolerising cell and the therapeutic cell are derived from the  
10 same embryonic stem cell (and therefore have the same antigenic characteristics). It will be appreciated that the cells in question are foreign to the patient to be treated since if they are from the patient to be treated, there is no need for pretolerisation.

15 Typically, the tolerising cell is a cell which has good expression of MHC molecules, since these molecules are the principle antigenic determinants relevant to transplant rejection. Good expression of other antigens that may be relevant to transplant rejection is also desirable. The tolerising cell may be the same as the therapeutic cell; however, it may also be a precursor of the  
20 therapeutic cell ie a cell which is capable of differentiating into the therapeutic cell and which is already committed to differentiate into the therapeutic cell. Thus, the tolerising cell is not a pluripotent cell (ie one which can differentiate into any cell) since such cells may spontaneously form teratomas and are not suitable for administration to a patient. If the tolerising cell is not the same as  
25 the therapeutic cell it is preferred if it is a precursor cell which is one or two or three or more stages less differentiated (on the same differentiation pathway) as the therapeutic cell.

In connection with the use of adult stem cells as the therapeutic cells, it may  
30 be advantageous to use cells derived from the peripheral blood of the donor as

the tolerising cells. Suitable cells include peripheral blood leukocytes which have good expression of MHC antigens. Alternatively, stem cells isolated from peripheral blood (eg a monocyte-derived subset; Zhao *et al* (2003) *Proc. Natl. Acad. Sci. USA* 100, 2426-2431) may be differentiated using, for example, EGF to give an epithelial phenotype, and used as tolerising cells.

The therapeutic cell is any suitable therapeutic cell. It may be a cell which is the same as the cell type which is damaged or diseased in the patient or one which is able to generate tissue which is damaged or diseased in the patient. Preferably, the cell is a precursor of the cell or tissue to be replaced or repaired, which is able to differentiate into the cell or tissue which is to be replaced or repaired. The cell is one which is already committed to differentiate into the cell or tissue to be replaced or repaired, and is not pluripotent (since such cells may cause the production of teratomas as discussed above).

The therapeutic cell (and therefore consequently the tolerising cell) is chosen by reference to the disease or disorder to be treated. Thus, typically, the therapeutic cell is, or is able to differentiate into, the cell or tissue which is to be regenerated in treating the disease or disorder. For example, in relation to precursor cells, in the case of diabetes, the precursor cell is one which is able to differentiate into an insulin-producing cell; in the case of congestive heart failure, the precursor cell is one which is able to differentiate into heart muscle cells; in the case of Parkinson's disease, the precursor cell is one which is able to differentiate into a suitable nerve cell; and so on. Suitable precursor cells which are able to differentiate into a type of cell or tissue which is used to replace the function of a failed or damaged cell or tissue in a degenerative disease or disorder are known in the art. Figure 10 describes stem cell lineages for human pluripotent stem cells (hPSCs), and suitable precursor cells, including stem cells (but not pluripotent stem cells as discussed above),

may be selected by reference to this figure. It will be appreciated that the precursor cell may be at one stage removed from the stage of differentiation where the function of the failed or damaged cell or tissue is expressed, or it may be two or three or four or more stages removed, but in each case the precursor cell is able to differentiate into the functional cell or tissue relevant to the disease to be treated.

The tolerising cells and the therapeutic cells may be, or be derived from, allogeneic adult stem cells (also called somatic stem cells). Typically, however, the tolerising cells and the therapeutic cells are derived from (but are not) embryonic stem cells, which are allogeneic. Human embryonic stem cells are typically from supernumerary embryos donated by couples who have benefited from successful *in vitro* fertilisation (IVF) cycles and have frozen embryos that are not required in the context of the IVF treatment. Protocols for the derivation of human stem cells are well known in the art, some of which are described in US Patent No 6, 280, 718 B1, incorporated herein by reference.

Derivatives of human embryonic stem cells (eg those which are lineage-specific stem cells) are functionally and physiologically similar, and sometimes identical to, somatic stem cells which all humans have and which provide us with a limited ability to repair and regenerate certain tissues. These include:-

(i) Hematopoietic stem cells that give rise to all the types of blood cells: red blood cells, B lymphocytes, T lymphocytes, natural killer cells, neutrophils, basophils, eosinophils, monocytes, macrophages, and platelets.

(ii) Bone marrow stromal cells (mesenchymal stem cells) that give rise to a variety of cell types; bone cells (osteocytes), cartilage cells (chondrocytes), fat

cells (adipocytes), and other kinds of connective tissue cells such as those in tendons.

(iii) Neural stem cells in the brain that give rise to its three major cell types: nerve cells (neurons) and two categories of non-neuronal cells - astrocytes and oligodendrocytes.

(iv) Epithelial stem cells in the lining of the digestive tract occur in deep crypts and that give rise to several cell types: absorptive cells, goblet cells, Paneth cells, and enteroendocrine cells.

(v) Skin stem cells occur in the basal layer of the epidermis and at the base of hair follicles. These epidermal stem cells give rise to keratinocytes, which migrate to the surface of the skin and form a protective layer. The follicular stem cells can give rise to both the hair follicle and to the epidermis.

Methods of differentiating pluripotent embryonic stem cells in to lineage-specific stem cells are known in the art. For example, US Patent No 6,280,718 B1 to Kaufman and Thomson, incorporated herein by reference, describes a method of obtaining human haematopoietic stem cells from a culture of human pluripotent embryonic stem cells. US Patent No 6,458,589 B1 to Rambhatla and Carpenter, herein incorporated by reference, describes methods for producing hepatocyte lineage cells from pluripotent stem cells. Pfendler & Kawase (2003) *Obstetrical & Gynecological Survey* 58, 197-208 review other methods of differentiating embryonic stem cells into dopamine-producing neurons, myelin-producing oligodendrocytes, insulin-producing cells, cardiomyocytes and so on.

Although it is preferred if the cell type to be transplanted itself or a precursor thereof is used as the tolerising cell, an antigen found thereon may also be

used, particularly if the antigen is a major antigen leading to histoincompatibility. Thus, MHC molecules which match the MHC antigens on the therapeutic cell may be used for tolerisation. The MHC molecules may, for instance, be used on a suitable synthetic molecular scaffold.

5

It will be appreciated that it is convenient in the practice of the invention for a "master" embryonic stem cell to be kept from which it is possible to produce suitable tolerising cells and therapeutic cells. It may be particularly convenient to use the earliest cell derived from an embryonic stem cell but  
10 which is committed to a particular path of differentiation (to the cell or tissue to be repaired or replaced) as the tolerising cell, and a later cell from the same differentiation pathway as the therapeutic cell. In this way, while the patient is being tolerised (using the earlier cell), cells suitable as the therapeutic cells are being produced (both derived from the same master embryonic stem cell and  
15 having common antigenic characteristics).

It will be appreciated that the tolerising cell or the therapeutic cell or both may be natural cells or may be genetically engineered cells. The tolerising cells, for example, may be genetically engineered to be more immunogenic than  
20 natural cells and so be more efficient at tolerisation, for example by overexpression of MHC antigens. The therapeutic cells may be genetically engineered to enhance their therapeutic properties, for example, cells which are able to regenerate islets of Langerhans may be genetically engineered to better produce insulin.

25

It is appreciated that to induce tolerance to an antigen, a derivative of the antigen may be administered to the patient, and not the antigen itself. By "derivative" of an antigen we include any portion of the antigen which can be presented by a class I or a class II MHC molecule for example on an antigen  
30 presenting cell (APC), and which induces tolerance to the antigen itself. For

example, a suitable portion of an antigen is a proteolytic digest of an MHC Class II molecule from the donor. Typically the derivative of the antigen is also recognised by a T cell when presented, for example via a T cell receptor.

- 5 When the antigen is a protein, a derivative of the antigen is typically a peptide fragment of the antigen consisting of a contiguous sequence of amino acids of the antigen capable of MHC binding. Preferably, the fragment is between 6 and 100 amino acids in length. More preferably, the fragment is between 6 and 50 amino acids in length. Most preferably, the fragment is six, or seven,  
10 or eight, or nine, or ten, or eleven, or twelve, or thirteen, or fourteen, or fifteen, or sixteen, or seventeen, or eighteen, or nineteen, or twenty, or twenty-one, or twenty-two, or twenty-three, or twenty-four or twenty-five amino acids in length.
- 15 A derivative of the antigen may include a fusion of the antigen, or a fusion of a fragment of the antigen, to another compound, and which can be recognised by either a class I or a class II MHC molecule when presented, and which induces tolerance to the antigen itself. Typically, the fusion is one which can be processed by an APC so as to present a portion which is able to induce  
20 tolerance to the antigen itself.

Unless the context indicates otherwise, wherever the term "antigen" is used in the context of an antigen, a derivative as herein defined is included.

- 25 The agent which raises the effective cAMP concentration in a monocyte cell may do so in several distinct but related biochemical ways. Thus, the agent may be one which increases the production of cAMP, for example by the stimulation of receptors which are linked to the production of cAMP. Such agents include prostaglandins and agonists thereof which are described in  
30 more detail below. Cholera toxin can also be used to increase cAMP levels



intracellularly as has been described in Braun *et al* (1999) *J. Exp. Med.* 189, 541-552 and there is also evidence that it may increase antigen transport across the epithelium which may be desirable. Similarly,  $\beta$ -adrenergic agents, which raise cAMP levels within a cell *via* the  $\beta$ -adrenergic receptor, may be used. Such  $\beta$ -adrenergic agents are well known in the art, such as in the treatment of asthma. Suitable  $\beta$ -adrenergic agents include isoproterenol.

The agent may be one which inhibits the breakdown of cAMP and thus may be a cAMP phosphodiesterase inhibitor, which are described in more detail below. The agent may be one which inhibits the export of cAMP from the cell. Export of cAMP from the cell is via a specific transporter (typically the multidrug resistance protein, MRP-4) which may be blocked with, for example, probenecid (a drug currently used for gout) or progesterone or agonists or antagonists thereof, such as medroxyprogesterone acetate or RU 486, which also appears to have an inhibitory effect on the cAMP transporter.

The agent may also be a compound which mimics the effects of cAMP in the cell in relation to generating a pro-tolerant state but which may be less susceptible to degradation or export. Such compounds, when present in the cell can be considered to raise the effective cAMP concentration. Such compounds include Sp-adenosine 3',5'-cyclic monophosphorothioate and 8-bromoadenosine 3',5'-cyclic monophosphate and dibutyryl cAMP. That sufficient of these compounds have been administered may be assessed by determining that there has been an elevation in IL-10 expression in monocyte cells. Preferably, the agent when used at a concentration which gives a maximal response elevates IL-10 expression at least 1.2-fold, or 1.5-fold, or 2-fold, or 5-fold, or 10-fold. Typically, from around 1 to 100  $\mu$ mol of the cAMP analogues may be administered to the patient.

Forskolin is 7 $\beta$ -Acetoxy-8,13-epoxy-1 $\alpha$ ,6 $\beta$ ,9 $\alpha$ -trihydroxylabd-14-en-11-one  
7 $\beta$ -Acetoxy-1 $\alpha$ ,6 $\beta$ ,9 $\alpha$ -trihydroxy-8,13-epoxy-labd-14-en-11-one. It is also  
called Coleonol and Colforsin and has a  $M_r$  of 410. It is a cell-permeable  
diterpenoid that possesses anti-hypertensive, positive inotropic and adenylyl  
cyclase activating properties. Many of its biological effects are due to its  
activation of adenylyl cyclase and the resulting increase in intracellular  
cAMP concentration. Forskolin affects calcium currents and inhibits MAP  
kinase. Colforsin is used as daropate (see *Ann Thoracic Surgery* (2001) 71,  
1931-1938). It may be administered as the hydrochloride to ensure water  
solubility but it may also be used as the free base which may be able to more  
readily penetrate cell membranes.

Sp-Adenosine 3',5'-cyclic monophosphorothioate (SpcAMP) has a  $M_r$  of 446  
and is the Sp-diastereomer of adenosine-3',5'-cyclic monophosphothioate. It  
is a potent, membrane-permeable activator of cAMP dependent protein kinase  
I and II that mimics the effects of cAMP as a second messenger in numerous  
systems while being resistant to cyclic nucleotide phosphodiesterases. It  
exhibits greater specificity and affinity than forskolin and cAMP analogues  
such as dibutyl-cAMP.

8-Bromoadenosine 3',5'-cyclic monophosphate (8-BrcAMP) has a  $M_r$  of 430.  
It is a cell-permeable cAMP analogue having greater resistance to hydrolysis  
by phosphodiesterases than cAMP. It activates protein kinase A.

Cholera toxin has a  $M_r$  of around 100,000. It is a toxin consisting of an A  
subunit (27 kDa) surrounded by five B subunits (approximately 12 kDa each),  
which attach the toxin to ganglioside GM1 on the cell surface. The A subunit  
catalyzes ADP-ribosylation of the  $\alpha$ -subunit of the stimulatory G protein  
(Gas) reducing GTPase activity and activating the  $\alpha$ -subunit. This activation

of G $\alpha$ s leads to an increase in the activity of adenylate cyclase resulting in increased levels of cAMP. It also ADP-ribosylates transducin in the eye rod outer segments, inactivating its GTPase activity. Cholera toxin has also been reported to ADP-ribosylate tubulin. It has been shown to be a potent mucosal vaccine adjuvant, inducing T helper cell type 2 responses by inhibiting the production of interleukin-12 (Braun *et al* (1999) *supra*). Although fragments of cholera toxin which are able to increase cAMP levels in monocytes may be used, it is preferred that complete cholera toxin is used.

Since cholera toxin may, under some conditions, induce anaphylaxis (oversensitization), it is less preferred.

---

It is likely that SpcAMP and 8-BrcAMP, agents such as rolipram and possibly forskolin, inhibit the cAMP export pump and this may contribute to their ability for raising the effective cAMP concentration.

It is convenient to measure the effective cAMP concentration in monocyte cells (ie by assessing the effect of the agent on monocyte cells). A preferred monocyte cell is the well known human monocyte cell line U937. It will be appreciated that the agents will also raise the effective cAMP concentration in other monocyte and monocyte-related cells such as macrophages, and that the utility in the context of the invention may be due to the effect on these cells. As noted above, whether or not there is a sufficient amount of cAMP analogues can be determined by measuring IL-10 in monocyte cells. Preferably, the agent when used at a concentration which gives a maximal response raises the cAMP concentration at least 1.2-fold, or 1.5-fold, or 2-fold, or 5-fold, or 10-fold.

Figure 11 shows diagrammatically various places of intervention in or on a cell which lead to raising cAMP levels.

It is preferred that the agent which raises the effective cAMP concentration in a monocyte cell is a prostaglandin.

- 5 It is preferred for this and all other aspects of the invention that the prostaglandin or agonist thereof stimulates cAMP production in a monocyte.

The prostaglandin or agonist thereof may be any suitable prostaglandin or agonist thereof that stimulates cAMP production in a monocyte, and which  
10 particularly in the presence of GMCSF causes monocytes to express IL-10. Prostaglandins or agonists thereof that are suitable for use in the present invention may readily be determined by a person of skill in the art. Methods for assessing cAMP production in monocytes may be found in Burzyn *et al.*, (2000) and in Example 3, and methods for detecting IL-10 expression in and  
15 release from monocytes include those in Examples 1 and 3.

By "prostaglandin or agonist" we mean any compound which acts as a prostaglandin agonist on a prostaglandin receptor. The prostaglandin agonist may be, but need not be, a prostanoid. Typically, the prostaglandin or agonist  
20 is one which binds the EP2 or EP4 receptor. The prostaglandin may be a PGE, a PGD or a PGI, or an agonist thereof. Preferably, the prostaglandin is a PGE or an agonist thereof. It is appreciated that PGI may be too unstable to be useful as a pharmacological agent, however PGI<sub>2</sub> and stable analogues of PGI may be suitable. Preferably, the prostaglandin is not a PGF or an agonist  
25 thereof.

It is preferred that the prostaglandin or agonist thereof is PGE<sub>2</sub> or a synthetic analogue thereof. Synthetic analogues include those modified at position 15 or 16 by the addition of a methyl group or those where the hydroxyl has been  
30 transposed from position 15 to position 16. Preferred examples of analogues

of prostaglandin include Butaprost (an EP2 receptor agonist) and 11-deoxy PGE1 (an EP4 receptor agonist) and 19-hydroxy PGE. For the avoidance of doubt, the term "prostaglandin" includes naturally-occurring prostaglandins as well as synthetic prostaglandin analogues.

5

Suitable prostaglandins or agonists thereof include dinoprostone (sold as Propess by Ferring in Europe and Forest in the USA; sold as Prostin E2 by Pharmacia), gemeprost (sold by Farillon), misoprostol (which is sold as Cytotec by Searle and Pharmacia), alprostadil (which is sold as Caverject by Pharmacia and Viridal by Schwarz and MUSE by AstraZeneca) and limaprost.

10

~~Misoprostol is a PGE analogue which has EP2 and EP3 agonist effects. Its chemical structure is ( $\pm$ ) methyl 11 $\alpha$ , 16-dihydroxy-16-methyl-9-oxoprost-13-enoate.~~

15

An example of a non-prostanoid compound which acts as a prostaglandin agonist is AH23848, an EP4 receptor agonist.

EP2 agonists which may be useful in the practise of the invention include AH13205.

20

Suitable prostaglandins also include 19-hydroxy PGE1 and 19-hydroxy PGE2. Prostaglandin E agonists are described in EP 1 097 922 and EP 1 114 816, incorporated herein by reference.

25

Suitable prostaglandins or agonists thereof may also include any of the 19-hydroxy prostaglandin analogues described in US Patent No. 4,127,612, incorporated herein by reference.

It is preferred that the prostaglandin is prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) or 19-hydroxy PGE. Prostaglandins and agonists thereof, including PGE<sub>2</sub>, are commercially available, for example from Pharmacia and Upjohn as Prostin E2.

5 The inventors further believes that it may be beneficial to use a phosphodiesterase (PDE) inhibitor either alone or with other agents which raise the effective cAMP concentration in a monocyte cell. The principal receptors for prostaglandin E2 (PGE<sub>2</sub>) are the EP2 and EP4 sub-types; however, other receptor sub-types exist (namely EP1 and EP3). EP2 and EP4  
10 receptors couple with adenylyclase and use elevated cAMP as the messenger system. The levels of cAMP in tissue are governed both by its synthesis and by its catabolism by PDEs which can be blocked by specific PDE inhibitors. Thus, the inventor believes that the effect of a prostaglandin or agonist thereof (such as PGE) acting on its EP2 and EP4 receptors is to stimulate cAMP, and  
15 the addition of the PDE inhibitor provides a synergistic action on monocytes and macrophages resulting in a reduction in the immune and/or inflammatory response which is greater than the effect of the sum of the same amount of the prostaglandin or agonist thereof, or PDE inhibitor administered alone.

20 Moreover, the inventors have previously found that the combination of a prostaglandin and a PDE inhibitor markedly stimulate IL-10 and inhibit IL-12 expression in, and secretion from, cells of the immune system, resulting in a tolerising environment.

25 Thus in an embodiment, the composition may further comprise a PDE inhibitor.

The PDE inhibitor may be any suitable PDE inhibitor. Preferably, the PDE inhibitor is one which inhibits a PDE which is active in cAMP breakdown.

30 The PDEs which are known to be active in cAMP breakdown are those of the

types IV, VII and VIII. Preferably, the PDE inhibitors are selective for type IV or VII or VIII.

Most preferably, the PDE inhibitors are selective for type IV PDE. By  
5 "selective" we mean that the inhibitor inhibits the particular type of PDE inhibitor for which it is selective, more potently than another type. Preferably, the type IV selective inhibitor is at least 2 times more potent an inhibitor of type IV PDE than another PDE type. More preferably, the type IV selective inhibitor is at least 5 times, 10 times, 20 times, 30, times 40 times, 50 times,  
10 100 times, 200 times, 500 times or 1000 times more potent an inhibitor of type IV PDE than another PDE type.

Typically, the selective inhibitor is around 5 to 50 times more potent an  
inhibitor of the selected PDE type than another PDE type. Typically, the  
15 selective inhibitor is 5 to 50 times more potent an inhibitor of the selected PDE type than an inhibitor that is considered to be non-selective such as theophylline. Thus, theophylline is 30 times less effective than rolipram.

Preferably, selective inhibition is determined by a comparison of  $IC_{50}$  levels  
20 (Dousa (1999) *Kidney International* 55: 29-62).

Non-specific PDE inhibitors include caffeine, theophylline, 3-isobutyl-1-methylxanthine (IBMX) and pentoxifylline (3,7-dihydro-3,7-dimethyl-1-(5-oxohexyl)-1H-purine-2,6-dione), although caffeine is not as active as the  
25 others and so is less preferred. The  $IC_{50}$  value for IBMX is 2-50  $\mu$ M.

US patent No. 6,127,378, incorporated herein by reference, discloses phenanthridines substituted in the 6 position that are described as selective PDE inhibitors (mainly of type IV), that may be suitable for use in the  
30 methods of the invention.

Specific (or selective) type IV PDE inhibitors include rolipram (4-[3-cyclopentyloxy-4-methoxyphenyl]-2-pyrrolidinone) and Ro-20-1724 (4-[3-butoxy-4-methoxybenzyl]-2-imidazolidinone). The  $IC_{50}$  for rolipram is  
5 800nM, and the  $IC_{50}$  for Ro-20-1724 is 2  $\mu$ M.

Another suitable PDE type IV selective inhibitor is denbufylline (1,3-di-n-butyl-7-(2-oxopropyl)-xanthine).

10 CP 80 633 (Hanifin *et al* (1996) *J. Invest. Dermatol.* 107, 51-56), CP 102 995 and CP 76 593 are also all potent type IV inhibitors (available from Central Research Division, Pfizer Inc, Groton, CT).

Other high affinity type IV selective PDE inhibitors include CPD 840, RP  
15 73401, and RS 33793 (Dousa, 1999). The high affinity type IV selective PDE inhibitors have a  $K_i$  of approximately 1 nM while the lower affinity inhibitors have a  $K_i$  of about 1  $\mu$ M.

The disclosures in Dousa (1999); Müller *et al* (1996, *Trends Pharmacol. Sci.*  
20 17: 294-298); Palfreyman & Souness (1996, *Prog Med Chem* 33: 1-52); Stafford & Feldman (1996, *Annual Reports in Medicinal Chemistry* (vol 31) pp 71-80; Ed. Bristol, Academic Press, NY, USA); and Teixeira *et al* (1997, *Trends Pharmacol. Sci.* 18: 164-171) relating to type IV PDE selective inhibitors are incorporated herein by reference.

25

Typically, when a type IV PDE-selective inhibitor is administered orally, around 1 to 30 mg is used. Thus, a typical oral dose of rolipram or denbufylline is 1 mg or 5 mg or 10 mg or 30 mg. When a non-selective PDE inhibitor is used, such as theophylline, and it is administered orally, the dose is  
30 between 5 and 50 mg, such as 5 or 10 or 20 or 30 or 40 or 50 mg.



When the composition includes progesterone, it is preferred if the dose of progesterone is sufficient to provide levels of between 100 nM and 50  $\mu$ M.

5 Preferred combinations are:

- PGE
- PGE+Rolipram
- PGE+probenecid
- 10 PGE+Rolipram+probenecid
- Forskolin
- Forskolin+Rolipram
- Forskolin+Rolipram+probenecid
- 8-Bromo cAMP+probenecid
- 15 8-Bromo cAMP+Rolipram+probenecid
- Sp-Adenosine 3,5-cyclic monophosphothioate (SpcAMP)
- SpcAMP+probenecid
- SpcAMP+Rolipram+probenecid
- Cholera toxin
- 20 Cholera toxin+probenecid

Preferably, these (and other agents which raise the effective cAMP concentration in a monocyte cell) are combined with GMCSF.

- 25 The inventors believe that these (and other) combinations may act synergistically to desirably raise the effect cAMP levels in monocyte cells. It will also be appreciated that by manipulating all the metabolic points for cAMP (see Figure 11), a lower dose of the components of the mixture would be possible in order to give the same effect compared to a single component
- 30 alone.

By "GMCSF" we include the gene product of the human GMCSF gene and naturally occurring variants thereof. The nucleotide and the amino acid sequence of human GMCSF is found in Genbank Accession No. NM\_000758, and in Figure 1. Some naturally occurring variants of GMCSF are also listed in NM\_000758. GMCSF is also known as colony stimulating factor 2 (CSF2).

The invention includes the use of derivatives of GMCSF that retain the biological activity of wild-type GMCSF, ie that stimulate the production of granulocytes and macrophages from their progenitor cells, and which in the presence of prostaglandin E cause monocytes to express IL-10.

By "derivative" of GMCSF we include a fragment, fusion or modification or analogue thereof, or a fusion or modification of a fragment thereof.

By "fragment" of GMCSF we mean any portion of the glycoprotein that stimulates the production of granulocytes and macrophages from their progenitor cells and which in the presence of prostaglandin E causes monocytes to express IL-10. Typically, the fragment has at least 30% of the activity of full length GMCSF. It is more preferred if the fragment has at least 50%, preferably at least 70% and more preferably at least 90% of the activity of full length GMCSF. Most preferably, the fragment has 100% or more of the activity of full length GMCSF.

The derivatives may be made using protein chemistry techniques for example using partial proteolysis (either exolytically or endolytically), or by *de novo* synthesis. Alternatively, the derivatives may be made by recombinant DNA technology. Suitable techniques for cloning, manipulation, modification and expression of nucleic acids, and purification of expressed proteins, are well known in the art and are described for example in Sambrook *et al* (2001)

"*Molecular Cloning, a Laboratory Manual*", 3<sup>rd</sup> edition, Sambrook *et al* (eds), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA, incorporated herein by reference.

- 5 The invention also includes modifications of full length GMCSF, or a fragment thereof, that stimulate the production of granulocytes and macrophages from their progenitor cells and which in the presence of prostaglandin E cause monocytes to express IL-10.
- 10 Such modifications include deglycosylating the glycoprotein either fully or partially. Other modifications include full length GMCSF, or a fragment thereof, having a different glycosylation pattern from that found in naturally occurring human GMCSF.
- 15 Other modifications of full length GMCSF, or a fragment thereof, include amino acid insertions, deletions and substitutions, either conservative or non-conservative, at one or more positions. Such modifications may be called analogues of GMCSF. By "conservative substitutions" is intended combinations such as Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr, Lys, Arg; and Phe, Tyr. Such modifications may be made using the methods of protein engineering and site-directed mutagenesis, as described in Sambrook *et al* 2001, *supra*. Preferably, the modified GMCSF or modified GMCSF fragment retains at least 30% of the activity of full length GMCSF. It is more preferred if the modified GMCSF or GMCSF derivative has at least 50%, preferably at least 70% and more preferably at least 90% of the activity of full length GMCSF. Most preferably, the modified GMCSF or modified GMCSF fragment has 100% or more of the activity of full length GMCSF.
- 25

The invention also includes the use of a fusion of full length GMCSF, or a fragment thereof, to another compound. Preferably, the fusion retains at least

30

30% of the activity of full length GMCSF. It is more preferred if the fusion has at least 50%, preferably at least 70% and more preferably at least 90% of the activity of full length GMCSF. Most preferably, the fusion has 100% or more of the activity of full length GMCSF.

5

GMCSF and analogues thereof are described in the following publications, each of which are incorporated herein by reference: US Patent No. 5,229,496 (Deeley *et al.*); US Patent No. 5,391,485 (Deeley *et al.*); US Patent No. 5,393,870 (Deeley *et al.*); US Patent No. 5,602,007 (Dunn *et al.*); Wong *et al.*,  
10 "Human GM-CSF: molecular cloning of the complementary DNA and purification of the natural and recombinant proteins", *Science* 228 (4701), 810-815 (1985); Lee *et al.*, "Isolation of cDNA for a human granulocyte-macrophage colony-stimulating factor by functional expression in mammalian cells", *Proc. Natl. Acad. Sci. U.S.A.* 82 (13), 4360-4364 (1985); Cantrell *et al.*,  
15 "Cloning, sequence, and expression of a human granulocyte/macrophage colony-stimulating factor", *Proc. Natl. Acad. Sci. U.S.A.* 82 (18), 6250-6254 (1985); and Miyatake *et al.*, "Structure of the chromosomal gene for granulocyte-macrophage colony stimulating factor: comparison of the mouse and human genes", *EMBO J.* 4 (10), 2561-2568 (1985).

20

While it is preferred that GMCSF is human GMCSF as defined above, by GMCSF we also include GMCSF from other species. However, it is appreciated that for applications in which GMCSF is administered to a subject, the GMCSF is preferably from the same species as the subject. Thus if the  
25 GMCSF is to be administered to a human subject, the GMCSF is preferably human GMCSF.

Suitable GMCSF for the practice of this invention can be obtained from Peptotech EC Ltd., 29 Margravine Road, London, W6 8LL, catalogue number  
30 300-03.

A preferred GMCSF for the practice of this invention is sargramostim, the proper name for yeast-derived recombinant human GMCSF, sold under the trade name Leukine<sup>®</sup> produced by Immunex, Inc. Leukine<sup>®</sup> is a recombinant  
5 human GMCSF produced in a *S. cerevisiae* expression system. Leukine<sup>®</sup> is a glycoprotein of 127 amino acids characterised by 3 primary molecular species having molecular masses of 19,500, 16,800 and 15,500 Daltons. The amino acid sequence of Leukine<sup>®</sup> differs from natural human GMCSF by a substitution of leucine at position 23, and the carbohydrate moiety may be  
10 different from the native protein. Leukine<sup>®</sup> is suitable for subcutaneous or intravenous administration (Leukine<sup>®</sup> Package Insert Approved Text, February 1998).

Unless the context indicates otherwise, wherever the term "GMCSF" is used, a  
15 derivative as herein defined is included.

In an embodiment, a monocyte-attracting chemotactic agent may also be used in aiding the production of a tolerising environment by attracting monocytes.

20 Suitable chemotactic agents for the practice of this invention include MIP-1 $\alpha$  and MCP-1, which can be obtained from Peprotech EC Ltd., 29 Margravine Road, London, W6 8LL, catalogue number 300-04. Other suitable chemotactic agents are described in US Patent No. 5,908,829 to Kelly, incorporated herein by reference.

25

Typically in the first, second and third aspects of the invention, the tolerising cell or an antigen found thereon or a derivative of said antigen and the agent which raises the effective cAMP concentration in a monocyte cell are administered simultaneously to the patient. More typically, they are all

present in the same composition (such as a pharmaceutical composition or formulation; see below). However, it is possible for the components to be administered separately, in which case it is desirable that the agent which raises the effective dAMP concentration in a monocyte cell are administered  
5 prior to administration of the tolerising cell or an antigen found thereon or a derivative of said antigen. Typically, if there is a time lag between administering the agent and the cells, it will be of the order of minutes.

Typically, in the third aspect of the invention, the therapeutic amount of the  
10 therapeutic cells are administered after tolerance has been achieved by the administration of the tolerising cells and the said agent. Thus, the patient is pretolerised to the cells prior to the therapeutic administration of therapeutic cells. Typically, the time between the pretolerisation regime and the therapeutic administration of therapeutic cells is of the order of 1 to 10 days.

15 The administration of the tolerising cells and the said agent typically is to a convenient site where the components can interact with the immune system and give rise to the induction of tolerance. Conveniently, a "tolerising" complex of the cells and the agent is used and administered to a mucous  
20 membrane which can be accessed non-invasively. Thus, suitable mucous membranes include those found in the mouth, vagina, anus, gastrointestinal tract and nose. Typically, therefore, the components are formulated as a buccal tablet, as a pessary, vaginal tablet or ring, or as a suppository or as a nasal spray.

25 Administration of the therapeutic amount of the precursor cell is directly or indirectly to the site where it is required in order regenerate failed or damaged cells or tissues. Typically, this is to the site of degeneration or damage or trauma and will vary depending on the disease or disorder to be treated.

The number of tolerising cells used may vary but would typically be around 100 to  $10^6$  cells. Sufficient of the agent which raises effective cAMP concentration in a monocyte cell is administered in order to produce a tolerising environment in the patient. Typically, around 2  $\mu\text{mol}$  of prostaglandin may be administered, around 50-100 ng GM-CSF and around 10  $\mu\text{mol}$  of a PDE inhibitor. When combinations are used, it is envisaged that lower amount of individual components will be required.

Sufficient therapeutic cells are administered to give a beneficial effect, such as initiation of repair or regeneration of the diseased or damaged cells or tissues. Typically around  $10^5$  to  $10^8$  therapeutic cells are administered, such as  $10^6$  or  $10^7$  cells.

How the cells are introduced to the site of disease or trauma varies. For example, for the treatment of diabetes, the "Edmonton protocol" may be used in which islet cells or immediate precursors thereof are injected into the portal vein of the liver in which organ they form effective, physiologically normal, glucose responsive and insulin-producing islets (see <http://www.diabetes.org.uk/islets/trans/edmonton.htm> for more details of the protocol). For the treatment of Parkinson's disease, cells may be injected into one of the accessible ventricles or portal veins from where they migrate to the correct site in the substantia nigra. For cardiac indications, and spinal chord injuries, cells may be injected directly to the site of damage to be repaired.

A further aspect of the invention provides a composition for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof comprising (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP

concentration in a monocyte cell, optionally, (c) granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof. As noted above, such a composition is useful for inducing tolerance. Conveniently, the composition is packaged and presented for use as a medicament, including as  
5 a medicament for human or veterinary use. Typically, the composition is packaged and presented for use in inducing tolerance to the therapeutic cell.

A still further aspect of the invention provides a therapeutic system for inducing tolerance to a therapeutic cell in a patient who is to be administered  
10 subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof comprising (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell, optionally, (c) granulocyte-macrophage  
15 colony stimulating factor (GMCSF) or a derivative thereof.

A yet still further aspect of the invention provides a kit of parts for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor  
20 thereof comprising (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell, optionally, (c) granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof.

25

The therapeutic system and kit of parts again are useful for inducing tolerance in a patient to the therapeutic cell. Optionally, the therapeutic system or kit of parts may additionally contain a therapeutic cell which is, or is able to differentiate into, the cell or tissue to be regenerated.

30



A pharmaceutical composition comprising a composition for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof comprising (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell, optionally, (c) granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof, and a pharmaceutically acceptable carrier is also included in the invention.

10

The carrier, diluent or excipient must be "acceptable" in the sense of being compatible with the composition of the invention and not deleterious to the recipients thereof. Typically, the carriers will be water or saline which will be sterile and pyrogen free. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985). The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as, or in addition to, the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), or solubilising agent(s).

Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

Further aspects of the invention include the following:

Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen and (b) an agent which raises the effective cAMP concentration in a monocyte cell, in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof.

Use of a tolerising cell sharing the same antigenic characteristics as a therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell wherein the patient is administered an agent which raises the effective cAMP concentration in a monocyte cell.

Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell wherein the patient is administered a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen.

Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and (c) GM-CSF, in the manufacture of a medicament for inducing tolerance to the therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the

therapeutic cell, and who is administered one or both of (a), (b) or (c) which is not found in the medicament as said.

Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as a therapeutic cell which therapeutic cell is, or is able to  
5 differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell, in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in  
10 need of transplantation of a therapeutic cell for cell or tissue regeneration.

~~Use of a tolerising cell sharing the same antigenic characteristics as a~~  
~~therapeutic cell which therapeutic cell is, or is able to differentiate into, the~~  
~~cell or tissue to be regenerated, or an antigen found thereon or a derivative of~~  
15 ~~said antigen in the manufacture of a medicament for reducing the risk of~~  
~~rejection of a transplant in a patient in need of transplantation of a therapeutic~~  
~~cell for cell or tissue regeneration wherein the patient is administered an agent~~  
~~which raises the effective cAMP concentration in a monocyte cell.~~

20 Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration wherein the patient is administered a tolerising cell sharing the same antigenic characteristics as the therapeutic cell which therapeutic cell  
25 is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen.

Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as a therapeutic cell which therapeutic cell is, or is able to  
30 differentiate into, the cell or tissue to be regenerated, or an antigen found

thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and (c) GMCSF in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration, and  
5 who is administered one or both of (a), (b) or (c) which is not found in the medicament as said.

Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which  
10 therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration, wherein the patient is subsequently administered a  
15 therapeutic amount of the said therapeutic cell.

Use of a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen  
20 found thereon or a derivative of said antigen in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration wherein the patient is administered an agent which raises the effective cAMP concentration in a monocyte cell and is subsequently administered a therapeutic amount of the said therapeutic cell.

25

Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration wherein the patient is administered a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered  
30 subsequently which therapeutic cell is, or is able to differentiate into, the cell

or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen and is subsequently administered a therapeutic amount of the said therapeutic cell.

- 5 Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and
- 10 (c) GMCSF or a derivative thereof in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration, and who is administered one or two of (a), (b) or (c) which is not found in the medicament as said, wherein the patient is subsequently administered a therapeutic amount of the said therapeutic cell.

15

- Use of a therapeutic amount of a therapeutic cell which is, or is able to differentiate into, a cell or tissue to be regenerated in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration, wherein the patient has previously been administered (a) a tolerising cell
- 20 sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen and (b) an agent which raises the effective cAMP concentration in a monocyte cell and, optionally, (c) GMCSF.

- The invention will now be described in more detail with the aid of the
- 25 following Figures and Examples.

### Figure 1

cDNA and amino acid sequence (Figures 1A and 1B, respectively) of human GMCSF, taken from GenBank Accession No. NM\_000758.

**Figure 2**

Figure 2 is a graph showing the effect of PGE and GMCSF on gene expression in U937 cells. Cells were treated for 4 hours with PGE2, with and without GMCSF, washed to remove the treatment, and incubated for a further 20 hours before the cells were pelleted and RNA extracted. The mRNA levels of CD14, CD80, CD86, BCL-2, BAX, COX-1 (cyclo-oxygenase 1), COX-2, PGES (prostaglandin synthase), EP2 (a prostaglandin receptor), EP4 (a prostaglandin receptor), PDE4B (a phosphodiesterase), IRAK-M, CIITA (MHC class II transactivator), MHC-II, IL-10 and granulysin (abbreviated to granlin), were measured. The graph indicates the percentage change in expression levels in the presence of GMCSF and PGE2.

**Figure 3**

Figure 3 is a graph showing the synergistic effect of PGE and GMCSF on the production of IL-10 mRNA in U937 cells, and that this phenotype is maintained 48 hours after removal of the treatment. Cells were treated for 4 hours with the agents indicated below the graph, washed to remove the treatment, and incubated for a further 48 hours before the cells were pelleted and RNA extracted. PGE2, E2 and E all refer to prostaglandin E2; GM refers to GMCSF; and M refers to MCSF.

**Figure 4**

Figure 4 is a graph showing the synergistic effect of PGE and GMCSF on the release of IL-10 protein in U937 cells, and that this phenotype is maintained after removal of the treatment. Cells were treated for 4 hours with the agents indicated below the graph, washed to remove the treatment, and incubated for a further 20 hours before the medium was assayed for IL-10. PGE refers to prostaglandin E2, and GM refers to GMCSF.

**Figure 5**

Expression of mRNA for cytokines IL-10 and IL-12 subunit p35. Experiment carried out on U937 cells (pro-monocytes) in the presence of Rolipram at 1  $\mu\text{g/ml}$  = 4  $\mu\text{M}$  and indomethacin 10  $\mu\text{M}$ . The indomethacin prevents  
5 prostaglandin synthesis from cells. Note that the effect of PGE+Rolipram is a marked stimulation of IL-10 and an inhibition of IL-12 both for unstimulated and IFN $\gamma$  stimulated cells. Vertical scale is a measure of mRNA compared to a control sample as measured by real-time quantitative PCR (Taqman).

**Figure 6**

Figure 6A is a graph showing the effect of PGE and Rolipram on the  
production of IL-10 mRNA in U937 cells. Figure 6B is a graph showing the  
effect of LPS, PGE and Rolipram on the production of IL-10 mRNA in U937  
cells. Figure 6C is a graph showing the effect of LPS, PGE and Rolipram on  
15 IL-10 release from U937 cells. Figure 6D is a graph showing the effect of PGE and Rolipram on IL-10 release from U937 cells.

**Figure 7**

A graph showing the effect of 19 hydroxy PGE1 and 19 hydroxy PGE2 on the  
20 stimulation of IL-10 in the presence and absence of rolipram.

**Figure 8**

A graph showing the effect of PGE1 and PGE2 on the stimulation of IL-10 in the  
presence and absence of rolipram.

25

**Figure 9**

A graph showing the effect of PGE and 19 hydroxy PGE on the production of  
phosphodiesterase IV b mRNA in the presence and absence of rolipram.

**Figure 10**

Stem cell lineages for human pluripotent stem cells.

**Figure 11**

- 5 Figure 11 is a diagram showing agents which control intracellular cAMP. Open arrows are effectively lowering intracellular cAMP levels. Solid arrow is stimulation. Combinations will be synergistic.

**Figure 12**

- 10 Figure 12 shows the relative efficacy of various agents in inducing IL-10 expression. See Example 4 for details.

**Figure 13**

- 15 Figure 13 shows the relative efficacy of various agents in inducing IL-10, expressed as a ratio of IL-10/TNF $\alpha$  mRNA expression. See Example 5 for details.

**Figure 14**

- 20 Figure 14 shows the relative efficacy of various agents and combinations of agents in inducing granulysin mRNA expression. See Example 6 for details.

**Figure 15**

- 25 Figure 15 shows that there is a synergistic effect between a prostaglandin (PGE2) and GMCSF and probenecid on the expression of IL-10.

**Example 1: Prostaglandin E / GMCSF synergism for inducing immunological tolerance**

- 30 There is growing evidence that prostaglandins of the E series are involved in immunological tolerance. This derives from their role in oral tolerance (the



ability of the immune system to distinguish pathogenic and comensal organisms), their ability to modulate cytokine ratios, and their huge concentrations in human seminal plasma where tolerance for the spermatozoon is essential.

5

Prostaglandins are produced at most mucosal surfaces of the body that have to accommodate beneficial or harmless bacteria and yet mount a response to pathogens. Newberry *et al* (1999) *Nature Medicine* 5, 900-906 have shown that 3A9 TCR $\alpha$   $-/-$  mice expressing a T cell receptor that specifically  
10 recognises egg-white lysosyme do not mount an inflammatory response to this antigen unless prostaglandin synthesis is inhibited, in that case by inhibiting the inducible cyclooxygenase isoform COX-2. With the source of prostaglandin removed, and with exposure to the specific antigen, these mice develop a pathology resembling inflammatory bowel disease (Newberry *et al*  
15 (1999) *supra*). These experiments confirm earlier studies showing that non-steroidal anti-inflammatory drugs such as indomethacin, which have a primary effect of inhibiting prostaglandin synthesis, break tolerance (Scheuer *et al* (1987) *Immunology* 104, 409-418; Louis *et al* (1996) *Immunology* 109, 21-26).

20

Monocytes of the normal lamina propria have a distinct phenotype since they express CD86 but not CD80. When an inflammatory condition persists (eg inflammatory bowel disease) the monocytes express CD80 (Rugtveit *et al* (1997) *Clin. Exp. Immunol.* 104, 409-418). The resident macrophages (CD80-  
25 ve CD86 +ve) are thus distinguished from the recently recruited macrophages which are CD80+ve, CD86+ve.

Monocytes are major sources of many immunological mediators, including prostaglandins and as such will alter the cytokine environment for antigen  
30 presentation. PGE has a major effect on cytokines relevant to tolerance,

stimulating the tolerogenic cytokine IL-10 (Strassmann *et al* (1994) *J. Exp. Med.* 180, 2365-2370) and inhibiting IL-12 (Kraan *et al* (1995) *J. Exp. Med.* 181, 775-779) which breaks tolerance. PGE will also have direct effects on the maturation of antigen-presenting dendritic cells, stimulating the production  
5 of cells that secrete increased IL-10 and diminished IL-12 (Kalinski *et al* (1997) *Adv. Exp. Med. Biol.* 417, 363-367).

A further indication of the importance of prostaglandins in ensuring essential tolerance is the very high (approximately millimolar) concentrations of both  
10 PGE and 19-hydroxy PGE in human seminal plasma. Clearly, immunological tolerance for spermatozoa entering the immunologically competent, and possibly infected, female genital tract is essential for the continuation of the species and levels of prostaglandin are such that many sub-epithelial, and even lymph-node cells will be affected. In this way, evolution has ensured  
15 immunological protection for the spermatozoa.

Previous experiments (Strassmann *et al* (1994) *supra*; Kraan *et al* (1995) *supra*) have required lipopolysaccharide (LPS) to be present for PGE to stimulate IL-10 production and in addition, the message for IL-10 was delayed  
20 by approximately 12 hours, both of these factors has been puzzling. The observations of the present invention suggest that LPS may have been stimulating the expression of GM-CSF, which may account for both the delay and the subsequent IL-10- expression.

25 We now show that the major prostaglandin effects on tolerance inducing monocytes may be mediated by a synergism between a prostaglandin and GM-CSF. The result of short term exposure to this combination results in a phenotype expressing greatly increased IL-10 but reduced levels of participants in antigen presentation such as CIITA and MHCII. Moreover,  
30 this change in phenotype is accompanied by enhanced expression of

granulysin. This molecule has anti-microbial properties (Krensky (2000) *Biochem. Pharmacol.* 59, 317-320) and is normally thought of as a product of activated T cells - mediating antiviral activity that lyses infected cells (Hata *et al* (2001) *Viral Immunol.* 14, 125-133; Ochoa *et al* (2001) *Nature Medicine* 7, 174-179; Smyth *et al* (2001) *J. Leukos. Biol.* 70, 18-29. Such an increase in innate defence molecules may compensate for the compromise of the adaptive immune system that necessarily accompanies tolerance induction. The phenotype is further characterised by a neutral effect on CD80 but a stimulation of CD86.

10

#### *Experimental Details*

U937 (human monocyte cell line) cells were grown in RPMI (PAA Laboratories) medium with 10% fetal calf serum added (PAA Laboratories). Cells were treated with prostaglandin E<sub>2</sub> at 10<sup>-6</sup> Molar with or without GM-CSF with at 5 ng/ml for 4 hours. The treatment was removed and cells were cultured for a further 20 hours. Cells were pelleted and the mRNA was extracted with Tri reagent (Sigma, Poole, UK). Total RNA was obtained by addition of chloroform and subsequent isopropanol precipitation. RNA was reverse transcribed with reverse transcriptase (Applied Biosystems) and random hexamers (Applied Biosystems). Probes and primers for amplification and detection of IL-10 and a number of other molecules were designed using Primer Express (Applied Biosystems) and are as follows:

#### IL-10 primers

CTACGGCGCTGTCATCGAT  
TGGAGCTTATTAAAGGCATTCTTCA

#### IL-10 probe

CTTCCCTGTGAAAACAAGAGCAAGGCC

30 BAX primers

45

CATGGAGCTGCAGAGGATGA  
CTGCCACTCGGAAAAGACCT

Bax Probe

TGCCGCCGTGGACACAGACTCC

5

BCL2 primers

CCGGGAGGCGACCGTAGT

GGGCTGCGCACCCCTTTC

BCL2 probe

10 CGCCGCGCAGGACCAGGA

CD80 primers

TCCACGTGACCAAGGAAGTG

CCAGCTCTTCAACAGAAACATTGT

15 CD80 Probe

AAGAAGTGGCAACGCTGTCCTGTGG

CD86 primers

CAGACCTGCCATGCCAATT

20 TTCCTGGTCCTGCCAAAATACTA

CD86 Probe

CAAACCTCTCAAAACCAAAGCCTGAGTGAGC

COX-1 primers

25 TGTTTCGGTGTCCAGTTCCAATA

ACCTTGAAGGAGTCAGGCATGAG

COX-1 Probe

CGCAACCGCATTGCCATGGAGT

30 COX-2 primers

46

GTGTTGACATCCAGATCACATTTGA

GAGAAGGCTTCCCAGCTTTTGTA

COX-2 Probe

TGACAGTCCACCAACTTACAATGCTGACTATGG

5

EP2 primers

GAC CGC TTA CCT GCA GCT GTA C

TGA AGT TGC AGG CGA GCA

EP2 Probe

10 CCA CCC TGC TGC TGC TTC TCA TTG TCT

---

EP4 primers

ACGCCGCCTACTCCTACATG

AGAGGACGGTGGCGAGAAT

15 EP4 Probe

ACG CGG GCT TCA GCT CCT TCC T

PDE4b primers

CCTTCAGTAGCACCGGAATCA

20 CAAACAAACACACAGGCATGTAGTT

PDE4b Probe

AGCCTGCAGCCGCTCCAGCC

Granulysin primers

25 CAGGGTGTGAAAGGCATCTCA

GGAGCATGGCTGCAAGGA

Granulysin Probe

CGGCTGCCCCACCATGGC

30 CD14 primers

47

GCGCTCCGAGATGCATGT  
AGCCCAGCGAACGACAGA  
CD14 Probe  
TCCAGCGCCCTGAACTCCCTCA

5

E synthase primers

CGGAGGCCCCCAGTATTG  
GGGTAGATGGTCTCCATGTCGTT

E synthase Probe

10 CGACCCCGACGTGGAACGCT

IRAKM primers

CCT GCC CTC GGA ATT TCT CT  
CTT TGC CCG CGT TGC A

15 IRAKM probe

CAC ACC GGC CTG CCA AAC AGA A

CIITA primers

GCTGTTGTGTGACATGGAAGGT  
20 RTGGGAGTCCTGGAAGACATACTG

CIITA Probe

CCGCGATATTGGCATAAGCCTCCCT

Class II primers

25 AGCCCAACGTCCTCATCTGT  
TCGAAGCCACGTGACATTGA

ClassII Probe

TCATCGACAAGTTCACCCACCAAGTG

Template was amplified in a Taqman 7700 machine for 40 cycles using FAM/TAMRA dyes on the probe. The Applied Biosystems Kit was used to amplify and detect ribosomal (18S) RNA as a control. After 40 cycles the Ct (related to cycle number at which signal appears) for the FAM and the 18S (VIC) were recorded and absolute relative quantitation was achieved using the formula  $2^{-\Delta\Delta C_t}$ .

The results of this experiment are shown in Figure 2 and show that there is a synergistic between a prostaglandin (PGE2) and GMCSF on the release of IL-10, CD-14, CD86, COX-2, and granulysin from cells of the immune system.

---

**Example 2: Prostaglandin E / GMCSF synergism for inducing IL-10**

Cells were cultured as described in Example 1 but after 4 hours medium was removed, cells were washed and the cells were cultured in medium alone for a further 48 hours. RNA was extracted from the cells as described in Example 1.

The results of this experiment are shown in Figure 3 and show that there is a synergistic effect between a prostaglandin (PGE2) and GMCSF on the expression of IL-10, and that this phenotype is maintained 48 hours after removal of the treatment.

**Example 3: Release of IL-10 from monocytes in response to PGE and GMCSF**

U937 cells were grown in RPMI (PAA Laboratories) medium with 10% foetal calf serum (PAA Laboratories) added. Cells were treated with prostaglandin E2 at  $10^{-6}$  Molar both with and without GMCSF at 5 ng/ml for 4 hours. The treatment was removed and cells were cultured for a further 20 hours.

Medium was removed and assayed for IL-10 using a matched monoclonal antibody pair (Pharmingen) or a commercial ELISA (R&D Systems, catalogue number D1000, Abingdon, Oxford). Figure 4 shows the release of IL-10 from monocytes in response to PGE and GMCSF.

5

To assay for cyclic AMP levels, wells in which cells are growing are treated with 0.01N hydrochloric acid to extract intracellular cAMP. This extract is neutralised to pH 6 and assayed for cyclic AMP in a competitive enzyme immunoassay (R&D Systems, catalogue number DE0450, Abingdon, Oxford).

10

**Example 4: Effect of the combination of PGE and rolipram on IL-10 and IL-12 production by U-937 (promonocyte) cells**

U 937 (human monocyte cell line) cells were grown in RPMI (PAA Laboratories) medium with 10% fetal calf serum added (PAA Laboratories). Cells were treated with prostaglandin E 2 at  $10^{-6}$  Molar or with Interferon- $\gamma$  at 10 ng/ml for 24 hours. Rolipram at 1  $\mu$ g/ml and indomethacin at 10  $\mu$ M was present in all wells. Cells were pelleted and the mRNA was extracted with Tri reagent (Sigma, Poole, UK). Total RNA was obtained by addition of chloroform and subsequent isopropanol precipitation. RNA was reverse transcribed with reverse transcriptase (Applied Biosystems) and random hexamers (Applied Biosystems). Probes and primers for IL-10 and IL-12 (p35) were designed using Primer Express (Applied Biosystems) and were as follows:

25

IL-12 p35 primers

CCACTCCAGACCCAGGAATG

TGTCTGGCCTTCTGGAGCAT

IL-12 probe

30 

TCCCATGCCTTCACCACTCCCAA



IL-10 primers

CTACGGCGCTGTCATCGAT

TGGAGCTTATTAAAGGCATTCTTCA

IL-10 probe

5 CTTCCTGTGAAAACAAGAGCAAGGCC

Template was amplified in a Taqman 7700 machine for 40 cycles using FAM/TAMRA dyes on the probe. The Applied Biosystems Kit was used to amplify and detect ribosomal (18S) RNA as a control. After 40 cycles the Ct  
10 (related to cycle number at which signal appears) for the FAM and the 18S (VIC) were recorded and absolute relative quantitation was achieved using the formula  $2^{-\Delta\Delta C_t}$ .

The results of this experiment are described in the legend to Figure 5. They  
15 show that there is a synergistic between a prostaglandin (PGE2) and a PDE inhibitor (rolipram) on the release of IL-10 from cells of the immune system and that there is a marked stimulation of IL-10 and inhibition of IL-12 in cells of the immune system when a prostaglandin (PGE2) and a PDE inhibitor (rolipram) are used in combination.

20

**Example 5: Stimulation of IL-10 production is achieved with or without LPS**

U 937 cells were grown in RPMI (PAA Laboratories) medium with 10% fetal  
25 calf serum added (PAA Laboratories).  $2 \times 10^6$  cells per flask were treated with prostaglandin E<sub>2</sub> at  $10^{-6}$  Molar or with Rolipram ( $4 \times 10^{-5}$ ) for 24 hours. Medium was removed at 20 hours and analysed by ELISA. A capture antibody (Pharmingen) was coated onto 96 well plates and culture medium was added each well. A standard curve was created with recombinant IL-10  
30 protein. After incubation and washing, a biotin labelled monoclonal antibody

(Pharmingen) was added and following incubation and washing, peroxidase labelled streptavidin was added. After washing a tetramethyl benzidine substrate was added and colour developed in proportion to IL-10 in the original sample/standard. Colour was read using a plate photometer (Labsystems, Multiskan). Mean concentrations (N=3) in controls with no lipopolysaccharide (LPS) were 38.2pg/ml and in the presence of LPS (100nM) they were 43.9 prostaglandin/ml.

After the incubation (20 hours), cells were pelleted and the mRNA was extracted with Tri-reagent (Sigma, Poole, UK). Total RNA was obtained by addition of chloroform and subsequent isopropanol precipitation. RNA was reverse transcribed with reverse transcriptase (Applied Biosystems) and random hexamers (Applied Biosystems). Probes and primers for IL-10 and IL-12 (p35) were designed using Primer Express (Applied Biosystems) and were as follows:

IL-12 p35 primers

CCACTCCAGACCCAGGAATG  
TGTCTGGCCTTCTGGAGCAT

IL-12 probe

TCCCATGCCTTCACCACTCCCAA

IL-10 primers

CTACGGCGCTGTCATCGAT  
TGGAGCTTATTAAAGGCATTCTTCA

IL-10 probe

CTTCCCTGTGAAAACAAGAGCAAGGCC

Template was amplified in a Taqman 7700 machine for 40 cycles using FAM/TAMRA dyes on the probe. The Applied Biosystems kit was used to amplify and detect ribosomal (18S) RNA (using VIC/TAMRA dyes) as an

internal control in the same reaction tube. After 40 cycles the Ct (related to cycle number at which signal appears) for the FAM and the 18S (VIC) were recorded and absolute relative quantitation was achieved using the formula  $2^{-\Delta\Delta C_t}$  where  $\Delta$  refers to the difference between the FAM and VIC signal related to an standard comparator included in each run.

### Example 6

The effect of PGE<sub>1</sub>, PGE<sub>2</sub>, 19 hydroxy PGE<sub>1</sub> and 19 hydroxy PGE<sub>2</sub> on the stimulation of IL-10 in the presence and absence of rolipram was investigated as described above in Example 5. IL-10 levels were measured using an ELISA assay (R&D Ltd, Oxford). Measurement was performed according to the manufacturer's instructions. Results are shown in Figures 7 and 8.

### Example 7

The mRNA for phosphodiesterase IV-b was measured as described in Example 5 above. mRNA was extracted after four hours of incubation. The concentration of the PGE was  $1 \times 10^{-6}$  and that of the 19-hydroxy PGE<sub>2</sub> was  $5 \times 10^{-6}$ . The following primers and Taqman probe were used for quantitation of PDE IV b mRNA.

Forward

CCTTCAGTAGCACCGGAATCA

Reverse

CAAACAAACACACAGGCATGTAGTT

Probe

AGCCTGCAGCCGCTCCAGCC

Results are shown in Figure 9. An increase in PDE activity follows both PGE and 19-hydroxy PGE application, which appears to be a direct negative feedback to reduce the effect of the stimulus. Use of a PGE and a type IV

selective PDE inhibitor increases PDE message levels even further, but then the synthesised phosphodiesterase is nullified by the presence of the inhibitor.

**Example 8: Relative efficacy of various agents which raise cAMP levels in monocyte cells in inducing IL-10**

*Experimental Details*

U937 (human monocyte cell line) cells were grown in RPMI (PAA Laboratories) medium with 10% fetal calf serum added (PAA Laboratories). Cells were treated with prostaglandin E2 at  $10^{-6}$  Molar, Rolipram  $10^{-6}$  Molar, Forskolin  $50 \times 10^{-6}$  Molar with or without GMCSF at 5 ng/ml for 48 hours. Cells were pelleted and the mRNA was extracted with Tri reagent (Sigma, Poole, UK). Total RNA was obtained by addition of chloroform and subsequent isopropanol precipitation. RNA was reverse transcribed with reverse transcriptase (Applied Biosystems) and random hexamers (Applied Biosystems). Probes and primers for amplification and detection of IL-10 were designed using Primer Express (Applied Biosystems) and are as follows:

IL-10 primers

CTACGGCGCTGTCATCGAT

TGGAGCTTATTAAAGGCATTCTTCA

IL-10 probe

CTTCCCTGTGAAAACAAGAGCAAGGCC

25

See Figure 12.

**Example 9: Relative efficacy of various agents which raise cAMP levels in monocyte cells in inducing IL-10 compared to TNF $\alpha$**

30

As for Example 4 but mRNA for TNF $\alpha$  is also included.

PMA ( $2 \times 10^{-7}$  M) was used as an alternative differentiating agent and although IL-10 was increased by PMA differentiation, TNF $\alpha$  (a pro-inflammatory and antitolerogenic agent) was also increased. Differentiation with Forskolin and GMCSF did not appreciably raise TNF $\alpha$ . Data is shown as the ratio of IL-10 mRNA/TNF $\alpha$  mRNA. P=PMA=Phorbol myristoyl acetate; F=Fsk=Forskolin, g=GMCSF, C=vehicle control.

10 TNF $\alpha$  Primers

GGAGAAGGGTGACCGACTCA

-----TGCCCAGACTCGGCAAAG-----

TNF $\alpha$  probe

15 CGCTGAGATCAATCGGCCCGACTA

See Figure 13.

20 Example 10: Relative efficacy of various agents which raise cAMP levels in monocyte cells in inducing granulysin

As for Example 4 but mRNA for granulysin was measured using the primers listed in Example 1 (see Figure 14).

25 G = GMCSF; FSK = Forskolin

Example 11: Prostaglandin E/GMCSF/probenecid synergism for inducing IL-10

Cells were cultured as described in Example 1 but after 20 hours medium was removed, cells were washed and RNA was extracted from the cells as described in Example 1.

- 5 The results of this experiment are shown in Figure 15 and show that there is a synergistic effect between a prostaglandin (PGE2) and GMCSF and probenecid on the expression of IL-10.

E = PGE2

10

**Example 12: Pre-tolerisation of patient undergoing stem cell treatment of diabetes mellitus**

- 15 Nestin-positive islet-derived precursor cells are isolated as described in Lechner *et al* (2002) *Biochem. Biophys. Res. Comm.* 293, 670-674. They are combined in a pessary with PGE2 and rolipram and/or GMCSF and the pessary inserted into the vagina of the female patient in order to tolerise the patient to the cells. Ten days later,  $10^6$ - $10^7$  of the nestin-positive islet-derived precursor cells are administered to the patient using the Edmonton protocol
- 20 protocol. In brief, the cells are injected into the hepatic portal vein of the patient from which they are taken to the liver where they form an insulin-producing islet.

**Example 13: Pre-tolerisation of patient undergoing stem cell treatment of**

25 **Parkinson's disease**

Human embryonic stem cells from an ethically approved donor source are differentiated into dopamine-producing neural cells by co-culture with PA6 cells, a stromal cell line derived from skull bone marrow. These are combined

56

with PGE2 and rolipram and/or GMCSF and formulated into a pessary. The patient is administered the pessary.

Once tolerance to the stem cells is achieved, the neural cells are introduced  
5 into one of the accessible ventricles or portal veins from where they migrate to  
the correct site in the substantia nigra of the patient and integrate.

10

**CLAIMS**

1. A method of inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof, the method comprising  
5 administering to the patient (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.
- 10 2. A method of reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration, the method comprising administering to the patient prior to the transplant (a) a tolerising cell sharing the same antigenic  
15 characteristics as the therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.
- 20 3. A method of treating a patient in need of cell or tissue regeneration the method comprising administering to the patient (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen  
25 found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell in an amount to induce tolerance to the said therapeutic cell, and subsequently administering to the patient (c) a therapeutic amount of the said therapeutic cell.



4. A method according to Claim 3 wherein in step (a) a cell is administered to the patient.
5. A method according to Claim 4 wherein the tolerising cell in step (a) and the therapeutic cell in step (c) are derived from the same parent embryonic stem cell.
6. A method according to any one of claims 1 to 5 wherein the patient is additionally administered granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof.
- ~~7. A method according to any one of the preceding claims wherein the patient is suffering from a degenerative disease or disorder.~~
8. A method according to Claim 7 wherein the degenerative disease or disorder is selected from the group consisting of diabetes, stroke, Parkinson's disease, ALS (Lou Gehrig's disease), spinal cord injury, heart attack, cardiac ischaemia, congestive heart failure, hepatitis, cirrhosis, cancer, immunodeficiency, osteoporosis, osteoarthritis, macular degeneration, burn, wounds, muscular dystrophy and multiple sclerosis.
9. A method according to any one of the preceding claims wherein (a) the tolerising cell or an antigen found thereon or a derivative of said antigen, and (b) the agent which raises the effective cAMP concentration in a monocyte cell are administered together.
10. A method according to Claim 9 wherein GMCSF is administered at the same time as (a) the tolerising cell, or an antigen found thereon or a

derivative of said antigen, and (b) the agent which raises the effective cAMP concentration in a monocyte cell.

- 5 11. A method according to any one of Claims 1 to 10 wherein (a) the tolerising cell or an antigen found thereon or a derivative of said antigen is administered after administration of (b) the agent which raises the effective cAMP concentration in a monocyte cell and, if used, the GMCSF or derivative thereof.
- 10 12. Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen and (b) an agent which raises the effective cAMP concentration in a monocyte cell, in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof.
- 15 13. Use of a tolerising cell sharing the same antigenic characteristics as a therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell wherein the patient is administered an agent which raises the effective cAMP concentration in a monocyte cell.
- 20 14. Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell wherein the patient is administered a tolerising cell sharing the same antigenic
- 25 30

characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen.

- 5 15. Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered therapeutically, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and (c) GM-CSF, in the manufacture of a medicament for inducing tolerance to the therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the therapeutic cell, and who is administered one or both of (a), (b) or (c) which is not found in the medicament as said.
- 10
- 
- 15 16. Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as a therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell, in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration.
- 20
- 25 17. Use of a tolerising cell sharing the same antigenic characteristics as a therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration wherein the patient is administered an agent which raises the effective cAMP concentration in a monocyte cell.
- 30

18. Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration wherein the patient is administered a tolerising cell sharing the same antigenic characteristics as the therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen.
19. Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as a therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and (c) GMCSF in the manufacture of a medicament for reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration, and who is administered one or both of (a), (b) or (c) which is not found in the medicament as said.
20. Use of a combination of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration, wherein the patient is subsequently administered a therapeutic amount of the said therapeutic cell.

21. Use of a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration wherein the patient is administered an agent which raises the effective cAMP concentration in a monocyte cell and is subsequently administered a therapeutic amount of the said therapeutic cell.
22. Use of an agent which raises the effective cAMP concentration in a monocyte cell in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration wherein the patient is administered a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen and is subsequently administered a therapeutic amount of the said therapeutic cell.
23. Use of any one or two of (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell and (c) GMCSF or a derivative thereof in the manufacture of a medicament for treating a patient in need of cell or tissue regeneration, and who is administered one or two of (a), (b) or (c) which is not found in the medicament as said, wherein the patient is

subsequently administered a therapeutic amount of the said therapeutic cell.

24. Use of a therapeutic amount of a therapeutic cell which is, or is able to  
5 differentiate into, a cell or tissue to be regenerated in the manufacture  
of a medicament for treating a patient in need of cell or tissue  
regeneration, wherein the patient has previously been administered (a) a  
tolerising cell sharing the same antigenic characteristics as the  
therapeutic cell, or an antigen found thereon or a derivative of said  
10 antigen and (b) an agent which raises the effective cAMP concentration  
in a monocyte cell and, optionally, (c) GMCSF.
25. A composition for inducing tolerance to a therapeutic cell in a patient  
who is to be administered subsequently a therapeutic amount of the said  
15 therapeutic cell or a precursor thereof comprising (a) a tolerising cell  
sharing the same antigenic characteristics as the therapeutic cell, or an  
antigen found thereon or a derivative of said antigen, (b) an agent  
which raises the effective cAMP concentration in a monocyte cell,  
optionally, (c) granulocyte-macrophage colony stimulating factor  
20 (GMCSF) or a derivative thereof.
26. A therapeutic system for inducing tolerance to a therapeutic cell in a  
patient who is to be administered subsequently a therapeutic amount of  
the said therapeutic cell or a precursor thereof comprising (a) a  
25 tolerising cell sharing the same antigenic characteristics as the  
therapeutic cell, or an antigen found thereon or a derivative of said  
antigen, (b) an agent which raises the effective cAMP concentration in  
a monocyte cell, optionally, (c) granulocyte-macrophage colony  
stimulating factor (GMCSF) or a derivative thereof.

27. A kit of parts for inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or a precursor thereof comprising (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell, optionally, (c) granulocyte-macrophage colony stimulating factor (GMCSF) or a derivative thereof.
28. A method according to any of Claims 1 to 11, a use according to any of Claims 11 to 24, a composition according to Claim 25, a therapeutic system according to Claim 26 or a kit of parts according to Claim 27 wherein the agent which raises the effective cAMP concentration in a monocyte cell is any one or more of a prostaglandin or agonist thereof, a  $\beta$ -adrenergic agent, a blocker of cAMP export from the cell, forskolin or a derivative thereof, a cAMP phosphodiesterase inhibitor, a cAMP analogue, or cholera toxin or a derivative or fragment thereof.
29. A method or use or composition or therapeutic system or kit of parts according to Claim 28 wherein the blocker of cAMP export from the cell is probenecid or progesterone.
30. A method or use or composition or therapeutic system or kit of parts according to Claim 28 wherein the cAMP analogue is Sp-adenosine 3',5'-cyclic monophosphorothioate or 8-bromoadenosine 3',5' cyclic monophosphate or dibutyryl cAMP.
31. A method or use or composition or therapeutic system or kit of parts according to Claim 28 wherein the prostaglandin or agonist thereof stimulates cAMP production in a monocyte.

32. A method or use or composition or therapeutic system or kit of parts according to Claim 28 or 31 wherein the prostaglandin or agonist thereof is any one of a prostaglandin E, such as prostaglandin E<sub>2</sub> or an analogue thereof, dinoprostone, gemeprost, misoprostol, alprostadil, limaprost, butaprost, 11-deoxy PGE<sub>1</sub>, AH23848, AH13205, or a 19-hydroxy PGE.
33. A method according to Claim 6 or a use according to any of Claims 15 or 19 or 23 or a composition according to Claim 25 or a therapeutic system according to Claim 26 or a kit of parts according to Claim 27 wherein the GMCSF, if present, is human GMCSF having the amino acid sequence as defined in Figure 1, or naturally occurring variants thereof.
34. A method or use or composition or therapeutic system or kit of parts according to Claim 33 wherein the GMCSF is sargramostim.
35. A method according to any of Claims 1 to 11 comprising administering a monocyte chemotactic agent to the patient.
36. A method according to Claim 35 wherein the monocyte chemotactic agent is MCP-1 or MIP-1 $\alpha$ .
37. A method according to any of Claims 1 to 11 comprising administering a PDE inhibitor to the patient.
38. A method or use or composition or therapeutic system or kit of parts according to Claim 28 wherein the PDE inhibitor is any one of 3-isobutyl-1-methylxanthine (IBMX), pentoxifylline (3,7-dihydro-3,7-



dimethyl-1-(5-oxohexyl)-1H-purine-2,6-dione), rolipram (4-[3-cyclopentyloxy-4-methoxyphenyl]-2-pyrrolidinone), CP80 633, CP102 995, CP76 593, Ro-20-1724 (4-[3-butoxy-4-methoxybenzyl]-2-imidazolidinone), theophylline, or denbufylline (1,3-di-n-butyl-7-(2-oxopropyl)-xanthine).

5

39. A method or use or composition or therapeutic system or kit of parts according to Claim 38 wherein the PDE inhibitor is selective for type IV PDE.

10

40. A method or use or composition or therapeutic system or kit of parts according to Claim 39 wherein the PDE inhibitor selective for type IV PDE is any one of rolipram (4-[3-cyclopentyloxy-4-methoxyphenyl]-2-pyrrolidinone), CP80 633, CP102 995, CP76 593, Ro-20-1724 (4-[3-butoxy-4-methoxybenzyl]-2-imidazolidinone), denbufylline (1,3-di-n-butyl-7-(2-oxopropyl)-xanthine), or CDP840, RP73401 or RS33793.

15

41. A pharmaceutical composition comprising the composition according to Claim 25 and a pharmaceutically acceptable carrier, diluent or excipient.

20

42. A composition according to Claim 25 for use in medicine.

43. A therapeutic system according to Claim 26 or a kit of parts according to Claim 27 further comprising a therapeutic cell which is, or is able to differentiate into, a cell or tissue to be regenerated.

25

44. Any novel method of pretolerising a patient to a cell as described herein.

30

67

45. Any novel method of pretolerising a patient to a cell as described herein, wherein the patient is subsequently administered a therapeutic amount of the cell or a precursor thereof in order to regenerate cells or tissue in the patient.

5

**ABSTRACT**

A method of inducing tolerance to a therapeutic cell in a patient who is to be administered subsequently a therapeutic amount of the said therapeutic cell or  
5 a precursor thereof, the method comprising administering to the patient (a) a tolerising cell sharing the same antigenic characteristics as the therapeutic cell, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.

10 A method of reducing the risk of rejection of a transplant in a patient in need of transplantation of a therapeutic cell for cell or tissue regeneration, the method comprising administering to the patient prior to the transplant (a) a  
15 tolerising cell sharing the same antigenic characteristics as the therapeutic cell which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, and (b) an agent which raises the effective cAMP concentration in a monocyte cell.

A method of treating a patient in need of cell or tissue regeneration the method comprising administering to the patient (a) a tolerising cell sharing the same  
20 antigenic characteristics as the therapeutic cell to be administered subsequently which therapeutic cell is, or is able to differentiate into, the cell or tissue to be regenerated, or an antigen found thereon or a derivative of said antigen, (b) an agent which raises the effective cAMP concentration in a monocyte cell in an amount to induce tolerance to the said therapeutic cell,  
25 and subsequently administering to the patient (c) a therapeutic amount of the said therapeutic cell.

Preferably, the agent which raises the effective cAMP concentration in a monocyte cell is a prostaglandin. Preferably it is used in combination with  
30 GMCSF or a derivative thereof.

Figure 2.

1/16

**FIGURE 1A**

Human GMCSF cDNA. GenBank Accession No. NM\_000758

1 gctggaggat gtggctgcag agcctgctgc tcttgggcac tgtggcctgc agcatctctg  
61 caccgccccg ctgcgccagc cccagcacgc agccctggga gcatgtgaat gccatccagg  
121 agggccggcg tctcctgaac ctgagtagag aactgctgc tgagatgaat gaaacagtag  
181 aagtcctctc agaaatgttt gacctccagg agccgacctg cctacagacc cgcctggagc  
241 tgtacaagca gggcctgcgg ggcagcctca ccaagctcaa gggcccttg accatgatgg  
301 ccagccacta caagcagcac tgccctccaa ccccggaac ttctgtgca acccagacta  
361 tcacctttga aagtttcaaa gagaacctga aggactttct gottgtcacc ccttttgact  
421 gctgggagcc agtccaggag tgagaccggc cagatgaggc tggccaagcc ggggagctgc  
481 tctctcatga aacaagagct agaaactcag gatgggtcacc ttggagggac caaggggtgg  
541 gccacagcca tgggtgggagt ggcctggacc tgccctgggc cacactgacc ctgatacagg  
601 catggcagaa gaatgggaat attttatact gacagaaatc agtaatatit atatatttat  
661 atttttaaaa tatttattta tttatttatt taagttcata ttccatattt attcaagatg  
721 ttttaccgta ataattatta ttaaaaatat gcttct

**FIGURE 1B**


Human GMCSF polypeptide. GenBank Accession No. NM\_000758

1 MWLQSLLLLG TVAC SISAPA RSPSPSTQFW EHVNAIQEAR RLLNLSRDTA AEMNETVEVI  
61 SEMFDLQEP T CLQTRLELYK QGLRGS LTKL KGPLTMMASH YKQHC PPTPE TSCATQTITF  
121 ESFKENLKDF LLVIPFDCWE PVQE

2/16

# FIGURE 2


Changes in gene expression in PGE+ GMCSF phenotype



3/16


**FIGURE 3**

**Message 48 hours after treatment**



4/16


FIGURE 4

**IL-10 release after 4hour  
treatment and 20 hours culture**

5/16

Figure 5A


13/3/00 IL-12 P35





6/16

Figure 5B



7/16

Figure 6A

mean  $\pm$  sem IL-10 message relative to control at  
20 hours





Figure 6B

IL-10 message in the presence of LPS



8/16

Figure 6C



**IL-10 release in presence of LPS**

Figure 6D

**IL-10 release no LPS**


**19-hydroxy PGE ± PDEI**

FIGURE 7

10/16




FIGURE 8

11/16

# Phosphodiesterase IVb mRNA (20 Hours)




FIGURE 9

# Stem-Cell Lineages





FIGURE 10

13/16

Figure 11



Agents controlling intracellular cAMP. Open arrows are effectively lowering intracellular cAMP levels. Solid arrow is stimulation. Combinations will be synergistic.



14/16

## Relative efficacy of inducing IL-10 expression




Figure 12





Figure 13

15/16



**Figure 14**

16/16




Figure 15

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**